

An Invariant for Varieties in Positive Characteristic

G. van der Geer and T. Katsura

ABSTRACT. We introduce an invariant of varieties in positive characteristic which generalizes the a -number of an abelian variety. We calculate it in some examples and discuss its meaning for moduli.

Dedicated to A.N. Parshin on the occasion of his 60th birthday

1. Introduction

Varieties in positive characteristic can differ markedly from varieties in characteristic zero. One aspect that makes this clear is Hodge theory. If X is a smooth variety in positive characteristic for which the Hodge-to-de Rham spectral sequence degenerates then the de Rham cohomology groups $H_{\text{dR}}^m(X)$ possess two filtrations, a decreasing filtration called the Hodge filtration F^\bullet and an increasing filtration called the conjugate filtration G_\bullet . The conjugate filtration is the analogue of the complex conjugate of the Hodge filtration in characteristic zero. But while in characteristic zero the Hodge filtration and the conjugate filtration are always transversal, this is no longer the case in positive characteristic and this gives rise to interesting new invariants of algebraic varieties in positive characteristic. Varieties in positive characteristic for which the Hodge filtration and the conjugate filtration are transversal are called ‘ordinary’ and resemble in some way smooth complex varieties, while varieties with non-transversal filtrations resemble singular complex varieties.

The relative position of the two filtrations on the de Rham cohomology is encoded by a double coset of a Weyl group and is rather complicated. We introduce an invariant which measures the position of the first step of the conjugate filtration in the Hodge filtration. If X is an abelian variety then this invariant coincides with the a -number as defined by Oort. The non-transversality is related to the cohomology of the sheaves $B_1\Omega^i$ introduced by Illusie.

In this paper after some preliminaries we define the a -number and show that this definition coincides with the definition of the a -number for abelian varieties and indicate the meaning of this number for the moduli of abelian varieties. This is related to work of Ogus on the order of singularity of the Hasse locus in the moduli of Calabi-Yau varieties. We then show how to calculate the a -number for Fermat varieties using the Poincaré residue map. After that we show an inequality for the

2000 *Mathematics Subject Classification.* 14K10.

Key words and phrases. a -number, Hodge filtration, conjugate filtration, Calabi-Yau.

a -number of Calabi-Yau threefolds and finish with a discussion of the a -number for a family of quintic Calabi-Yau varieties.

2. The Two Spectral Sequences

Let X be a smooth complete algebraic variety defined over an algebraically closed field k of characteristic $p > 0$. Let X' be the base-change of X with respect to the Frobenius homomorphism of k . The absolute Frobenius $F_{\text{abs}} : X \rightarrow X$ factors through the relative Frobenius $F_{\text{abs}} : X \xrightarrow{F} X' \rightarrow X$. There are two spectral sequences converging to the de Rham cohomology: the first is the Hodge spectral sequence

$$E_1^{i,j} = H^j(X, \Omega^i) \implies H_{\text{dR}}^{i+j}(X).$$

This spectral sequence arises from the filtration $\Omega^{\geq i}$ of the de Rham complex Ω^\bullet . The second spectral sequence is the so-called conjugate spectral sequence which comes from the the Leray spectral sequence for the relative Frobenius $F : X \rightarrow X'$, that yields a spectral sequence

$$E_2^{i,j} = H^i(X', \mathcal{H}(F_*\Omega_{X/k}^\bullet)) \implies H_{\text{dR}}^{i+j}(X/k).$$

But the Cartier operator yields an isomorphism of sheaves on X'

$$C^{-1} : \Omega_{X'/k}^i \xrightarrow{\sim} \mathcal{H}^i(F_*(\Omega_{X/k}^\bullet)),$$

so that we can rewrite this as

$$E_2^{i,j} = H^i(X', \mathcal{H}^j(F_*(\Omega_{X/k}^\bullet))) = H^i(X', \Omega_{X'/k}^j) \implies H_{\text{dR}}^*(X).$$

We assume that the Hodge-to-de Rham spectral sequence for X degenerates at the E_1 -level. This happens for example if $p > \dim(X)$ and X can be lifted to the Witt vectors $W_2(k)$, see [De-II]. Then the de Rham cohomology carries a filtration, the Hodge filtration

$$F^\bullet : H_{\text{dR}}^m = F^0 \supset F^1 \supset \dots \supset F^m,$$

with graded pieces

$$\text{gr}^i H_{\text{dR}}^m(X) = H^{m-i}(X, \Omega^i).$$

Moreover, then also the conjugate spectral sequence degenerates at the E_2 -level, leading to the so-called conjugate filtration

$$G_\bullet : (0) \subset G_0 \subset G_1 \subset \dots \subset G_m = H_{\text{dR}}^m(X)$$

with graded pieces

$$\text{gr}_i H_{\text{dR}}^m(X) = H^i(X', \Omega_{X'}^{m-i}).$$

Moreover, if $m = n := \dim(X)$ then we have a non-degenerate pairing \langle , \rangle on the de Rham cohomology

$$H_{\text{dR}}^n(X) \times H_{\text{dR}}^n(X) \rightarrow H_{\text{dR}}^{2n}(X) \cong k.$$

Note that for the conjugate spectral sequence we have

$$E_2^{n,0} = H^n(X', \mathcal{H}^0(\Omega_{X'/k}^\bullet)) = H^n(X', \mathcal{O}_{X'})$$

and since $F_{\text{abs}}^*(H^n(X, \mathcal{O}_X)) = H^n(X', \mathcal{O}_{X'})$ we see that $G_0 = F_{\text{abs}}^*(H^n(X, \mathcal{O}_X))$. In particular, the composition

$$H^n(X, \mathcal{O}_X) \xrightarrow{F^*} G_0 \rightarrow F^0/F^1 \cong H^n(X, \mathcal{O}_X)$$

is the Hasse-Witt map.

3. The Relative Position of Two Filtrations

Given two filtrations on the de Rham cohomology it is natural to compare them to obtain interesting information on the variety. In general, if one has two filtrations on a finite-dimensional k -vector space V

$$F^\bullet : V = F^0 \supset F^1 \supset \dots \supset F^m, \quad G_\bullet : G_0 \subset G_1 \subset \dots \subset G_m = V$$

such that $\text{rank}(G_i) = \text{rank}(F^{m-i})$ the relative position of two such filtrations is encoded by an element of a double coset of a Weyl group of the general linear group $\mathcal{G} = \text{GL}(V)$. Sometimes, the vector space V is provided with a non-degenerate pairing \langle , \rangle and then we can consider the group $\mathcal{G} = \text{GSp}(V)$ instead of $\text{GL}(V)$.

We fix a maximal torus T and a Borel subgroup B_0 containing T and let W be the Weyl group of \mathcal{G} . The (partial) flags F^\bullet and G_\bullet determine a parabolic subgroups P_F and P_G which are conjugate under W . Let $W_F = W_G$ be the Weyl group of F^\bullet . Then the relative position of F^\bullet and G_\bullet is given by an element of

$$w(F^\bullet, G_\bullet) \in W_F \backslash W / W_G.$$

To define it, recall that if \mathcal{B} denotes the variety of Borel subgroups of \mathcal{G} then we have a bijection $\phi : W \xrightarrow{\sim} \mathcal{G} \backslash \mathcal{B} \times \mathcal{B}$, given by $w \mapsto$ the orbit of (B, wBw^{-1}) . We choose Borel subgroups $B_F \subset P_F$ and $B_G \subset P_G$ and define $w(F^\bullet, G_\bullet)$ as the double coset containing (B_F, B_G) and this is independent of the choices of B_F and B_G . Note that the double coset $W_F \backslash W / W_G$ is in bijection with $P_G \backslash \mathcal{G} / P_F$, cf. [10].

DEFINITION 3.1. Let X be a smooth variety of dimension n in characteristic $p > 0$ for which the Hodge-to-de Rham spectral sequence degenerates and let m be an integer with $0 \leq m \leq 2n$. We define the pointer $w_m(X)$ of X as the element of $W_F \backslash W / W_G$ associated to the two filtrations $\{F^\bullet\}$ and $\{G_\bullet\}$ on $H_{\text{dR}}^m(X)$.

We can refine this considerably by choosing full filtrations refining the filtrations F^\bullet and G_\bullet .

$$V = \Phi^0 \supset \Phi^1 \supset \dots \supset \Phi^{\dim(V)}, \\ \Gamma_0 \subset \Gamma_1 \subset \dots \subset \Gamma_{\dim(V)} = V$$

that are compatible with the action of Frobenius in the following sense. Using crystalline cohomology we know that F/p^i acts on F^i/F^{i+1} and the induced filtration of Φ^\bullet should be stable under F/p^i . Moreover, the filtration Φ^\bullet induces via $G_i/G_{i-1} \cong F^{i-1}/F^i$ the filtration Γ_\bullet . Comparing the two filtrations gives the finer invariant. If we carry this out for principally polarized abelian varieties we retrieve the Ekedahl-Oort type of the abelian variety, cf. [14] where this invariant was introduced in a different way. We refer to [4] and to [3], where this will be worked out in detail for abelian varieties, K3-surfaces and Calabi-Yau's and leads to stratifications on moduli spaces (cf. also [10], [5]).

4. The a -Number of a Variety in Positive Characteristic

We are now interested in the case of the cohomology group $H_{\text{dR}}^m(X)$ with $m = n = \dim(X)$. We shall assume throughout that the Hodge-to-de Rham spectral sequence degenerates at the E_1 -level. Since the pointer $w_n(X)$ is a complicated invariant, we look first at the position of G_0 , the image of $H^n(X, \mathcal{O}_X)$ under the Frobenius operator in the Hodge filtration F^\bullet on $H_{\text{dR}}^n(X)$. We distill an invariant in the following way.

DEFINITION 4.1. The a -number of X is the maximum step in the Hodge filtration that contains $F_{\text{abs}}^*(H^n(X, O_X))$:

$$a(X) := \max\{j : F^j H_{dR}^n(X) \text{ contains } F_{\text{abs}}^*(H^n(X, O_X))\},$$

or equivalently,

$$a(X) = \max\{j : F^j \supseteq G_0\}.$$

Note that $0 \leq a(X) \leq n = \dim(X)$ and that $a(X) > 0$ if and only if the Hasse-Witt map vanishes. Of course, by going from $w_n(X)$ to $a(X)$ we lose a lot of information. If we wish to retain more information we can refine the a -number slightly by taking the a -vector $(a_0, a_1, \dots, a_m)(X)$ with

$$a_j(X) := \dim G_0 \cap F^j \quad j = 0, 1, \dots, m.$$

Sometimes, if $H^n(X, O_X)$ vanishes, we can still define an analogue of the a -number. Consider for example the case of a smooth cubic hypersurface in \mathbb{P}^8 . The Hodge numbers $h^{i,j}$ with $i + j = 7$ are

$$h^{0,7} = h^{7,0} = 0, \quad h^{1,6} = h^{6,1} = 0, \quad h^{2,5} = h^{5,2} = 1, \quad h^{3,4} = h^{4,3} = 360,$$

and we can define

$$a'(X) = \max\{j : F^j \supseteq G_2\}.$$

QUESTION 4.2. i) Can we give geometric interpretations of these numbers? ii) Is a variety with $a(X) = \dim(X)$ rigid?

5. Closed Differential Forms

Let X be a smooth projective variety of dimension n in characteristic $p > 0$ and let $F_{\text{abs}} : X \rightarrow X$ be the absolute Frobenius. We consider the direct image $F_{\text{abs}*} O_X$. As a sheaf of abelian groups it is just O_X , but its O_X -module structure is different: $f \circ g = f^p g$. One way to define the sheaf $B_1 \Omega_X^i$ is via the short exact sequence

$$0 \rightarrow \Omega_X^{i-1} \rightarrow F_{\text{abs}*} \Omega_X^{i-1} \xrightarrow{d} B_1 \Omega_X^i \rightarrow 0.$$

Here we view $B_1 \Omega_X^i$ as an O_X -module. It can also be viewed as a locally free subsheaf of $F_* \Omega_X^i$ on X' . Here F denotes the relative Frobenius. Moreover, we set $B_0 \Omega_X^i = 0$. Inductively we define

$$B_{j+1} \Omega_X^i = C^{-1}(B_j \Omega_X^i),$$

where $C : Z_1 \Omega_X^i \rightarrow \Omega_X^i$ is the Cartier operator. Similarly, we define

$$Z_0 \Omega_X^i = \Omega_X^i, \quad Z_1 \Omega_X^i = \Omega_{X, \text{d-closed}}^i$$

and inductively

$$Z_{j+1} \Omega_X^i = C^{-1}(Z_j \Omega_X^i) \quad \text{for } j \geq 1.$$

We can view these as locally free subsheaves of $F_*^j \Omega_X^i$ on $X^{(p^j)}$, the base change of X under the j -th power of relative Frobenius. The inverse Cartier operator gives rise to an isomorphism

$$C^{-j} : \Omega_{X^{(p^j)}}^i \xrightarrow{\cong} Z_j \Omega_X^i / B_j \Omega_X^i.$$

We have a perfect pairing

$$F_{\text{abs}*} \Omega_X^i \otimes F_{\text{abs}*} \Omega_X^{n-i} \rightarrow \Omega_X^n, \quad (\omega_1, \omega_2) \mapsto C(\omega_1 \wedge \omega_2).$$

On the other hand we have exact sequences

$$0 \rightarrow \Omega_X^i \rightarrow F_{\text{abs}*} \Omega_X^i \xrightarrow{d} B_1 \Omega_X^{i+1} \rightarrow 0$$

and

$$0 \rightarrow B_1 \Omega_X^{n-i} \rightarrow F_{\text{abs}*} \Omega_X^{n-i} \xrightarrow{C} \Omega_X^{n-i} \rightarrow 0.$$

This induces a perfect pairing

$$B_1 \Omega_X^{i+1} \otimes B_1 \Omega_X^{n-i} \rightarrow \Omega_X^n.$$

DEFINITION 5.1. (Illusie, Raynaud) We call the variety X *ordinary* if the cohomology groups $H^i(X, B_1 \Omega_X^j)$ vanish for $j \geq 1$ and all i .

This implies that all global forms are closed, just as in characteristic 0.

LEMMA 5.2. *If X is ordinary then $a(X) = 0$.*

PROOF. Consider the exact sequence

$$0 \rightarrow O_X \xrightarrow{F} O_X \xrightarrow{d} B_1 \Omega_X^1 \rightarrow 0.$$

In cohomology this gives

$$H^{n-1}(X, B_1 \Omega_X^1) \rightarrow H^n(X, O_X) \rightarrow H^n(X, O_X) \rightarrow H^n(X, B_1 \Omega_X^1),$$

from which it follows that the Hasse-Witt map has trivial kernel and thus that $G_0 \not\subseteq F^1(H_{\text{dR}}^n(X))$. \square

6. Abelian Varieties

Let X be an abelian variety of dimension g over an algebraically closed field k . We denote by $X[p]$ the kernel of multiplication by p . It is a group scheme of order p^{2g} . The classical a -number of X (cf. [13]) is defined by

$$a(X) := \dim_k \text{Hom}_k(\alpha_p, X).$$

Here α_p is the group scheme of order p that is the kernel of Frobenius acting on the additive group \mathbb{G}_a . Note that $\text{Hom}_k(\alpha_p, X)$, the space of group scheme homomorphisms of α_p to X , is in a natural way a vector space over k . If we let $A(X)$ be the maximal subgroup scheme of $X[p]$ annihilated by the operators F (Frobenius) and V (Verschiebung), then $A(X)$ is the union of the images of all group scheme homomorphisms $\alpha_p \rightarrow X$ and we have

$$a(X) = \log_p \text{ord} A(X).$$

We have $0 \leq a(X) \leq g$.

It is well-known that the contravariant Dieudonné module $D(X)$ of $X[p]$ can be identified with the first de Rham cohomology $H_{\text{dR}}^1(X)$ of X . The k -vector space $H_{\text{dR}}^1(X)$ then carries two operators F and V and the Dieudonné module of $A(X)$ coincides with the intersection $\ker(V) \cap \ker(F)$ on $D(X)$. If we identify the kernel of F with $H^0(X, \Omega_X^1) \subset H_{\text{dR}}^1(X)$ then the Dieudonné module of $A(X)$ may be identified with the kernel of V acting on $H^0(X, \Omega_X^1)$. We thus find

$$D(A(X)) \cong \ker(V : H^0(X, \Omega_X^1) \rightarrow H^0(X, \Omega_X^1)) \cong H^0(X, B_1 \Omega_X^1)$$

and we have

$$a(X) = \dim_k H^0(X, B_1 \Omega_X^1).$$

We also have the following relation:

$$F(H^1(X, O_X)) \cap H^0(X, \Omega_X^1) = H^0(X, B_1 \Omega_X^1). \tag{1}$$

This follows from the fact that $\text{Im}(F) = \text{Ker}(V)$ on $X[p]$. We now show that for abelian varieties the classical a -number and our a -number coincide.

PROPOSITION 6.1. *If X is an abelian variety of dimension g then the two notions of a -number coincide: if F^\bullet and G_\bullet are the Hodge and conjugate filtration on $H_{\text{dR}}^g(X)$ we have*

$$\dim_k \text{Hom}_k(\alpha_p, X) = \max\{j : G_0 \subset F^j\}.$$

PROOF. Recall that $H_{\text{dR}}^g(X) = \wedge^g H_{\text{dR}}^1(X)$ and if we write $H_{\text{dR}}^1(X) = V_1 \oplus V_2$ with $V_1 = H^0(X, \Omega_X^1)$ and V_2 a complementary subspace, then the Hodge filtration on $H_{\text{dR}}^g(X)$ is $F^r = \sum_{j=r}^g \wedge^j V_1 \otimes \wedge^{g-j} V_2$. We have $F(H^g(X, O_X)) = F(\wedge^g H^1(X, O_X)) = \wedge^g F(H^1(X, O_X))$. If we write $F(H^1(X, O_X)) = A \oplus B$ with $A = H^0(X, B_1 \Omega_X^1)$ and B a complementary space, then $\wedge^g(A \oplus B) = \wedge^a A \otimes \wedge^{g-a} B$ with $a = \dim(A)$. From this and (1) it is clear that $F(H^g(X, O_X))$ lies in F^a , but not in F^{a+1} . \square

We now show that the a -number has some meaning for the geometry of moduli spaces.

Let $T(a)$ be the locus inside the moduli space \mathcal{A}_g of principally polarized abelian varieties with a -number $\geq a$. Here \mathcal{A}_g is viewed as an algebraic stack, or we should add a sufficient level structure to our abelian varieties. Over \mathcal{A}_g the de Rham bundle $\mathcal{H}_{\text{dR}}^1$ possesses two subbundles of rank g , the Hodge bundle \mathbb{E} and the kernel \mathbb{F} of F . Then $T(a)$ is defined as the degeneracy locus where $\mathbb{E} \cap \mathbb{F}$ has rank at least a . It is known that $\dim T(a) = g(g+1)/2 - a(a+1)/2$, cf. [4], [3], [14].

PROPOSITION 6.2. *The locus $T(a)$ in \mathcal{A}_g is smooth outside $T(a+1)$ and the normal space to $T(a)$ at $[X]$ with $a(X) = a$ can be identified with $\text{Sym}^2(H^0(X, B_1 \Omega_X^1))$.*

PROOF. An infinitesimal deformation of the principally polarized abelian variety X is given by a symmetric $g \times g$ -matrix $T = (t_{ij})$ which can be interpreted as a symmetric endomorphism of $H^0(X, \Omega_X^1)$, cf. [9], p. 165. This deformation preserves the a -number of X if it keeps the kernel $H^0(X, B_1 \Omega_X^1)$ of V acting on $H^0(X, \Omega_X^1)$, that is, T is a symmetric endomorphism of this subspace. The principal polarization identifies this subspace with its dual, and this gives us the result. \square

PROPOSITION 6.3. *Let X be an abelian variety with $a(X) = g$. Then the multiplicity of the point $[X]$ on $T(1)$ is g .*

PROOF. If X is a principally polarized abelian variety with $a(X) = g$ we choose a base $\omega_1, \dots, \omega_g$ of $H^0(X, \Omega_X^1)$ and complete it to a basis of $H_{\text{dR}}^1(X)$ as η_1, \dots, η_g such that $\langle \omega_i, \eta_j \rangle = \delta_{ij}$. Then for a deformation with parameter a symmetric $g \times g$ -matrix (t_{ij}) we have for $i = 1, \dots, g$

$$\begin{aligned} F\eta_i &= \omega_1 + \sum t_{ij} \eta_j \\ \omega_i &= V(\sum t_{ij} \eta_i). \end{aligned}$$

Then we get

$$F(\eta_1 \wedge \dots \wedge \eta_g) = \omega_1 \wedge \dots \wedge \omega_g + \dots + \det(t_{ij}) \eta_1 \wedge \dots \wedge \eta_g.$$

But the equation of $T(1)$ is $\langle F(\eta_1 \wedge \dots \wedge \eta_g), \eta_1 \wedge \dots \wedge \eta_g \rangle = \det(t_{ij})$. \square

7. Cohomology of Hypersurfaces

Let X be an irreducible smooth projective hypersurface of degree d in \mathbb{P}^{n+1} given by an equation $f = 0$. Then the primitive cohomology $H_{\text{dR}}^n(X)_0$ in the middle dimension can be described by the Poincaré residue map as follows. Consider the rational differential forms on projective space

$$\Omega = \sum_{i=0}^{n+1} (-1)^i x_i dx_0 \wedge \dots \wedge d\hat{x}_i \wedge \dots \wedge dx_{n+1} \quad \text{and} \quad \Omega^* = \frac{\Omega}{x_0 \dots x_{n+1}}.$$

Note that Ω^* is a logarithmic form: in affine coordinates we have $\Omega^* = du_1/u_1 \wedge \dots \wedge du_{n+1}/u_{n+1}$.

We let \mathcal{L} be the k -vector space generated by the monomials $x^w = x_0^{w_0} \dots x_{n+1}^{w_{n+1}}$ with $\sum_{i=0}^{n+1} w_i \equiv 0 \pmod{d}$ and we let $\mathcal{L}' \subset \mathcal{L}$ be the subspace generated by the x^w with all $w_i \geq 1$, i.e., the elements in \mathcal{L}' are divisible by $x_0 \dots x_{n+1}$. For an element $x^w \in \mathcal{L}'$ we set $\gamma(w) = \sum w_i/d$.

On \mathcal{L} we have operators D_i defined by

$$D_i(x^w) = x_i \frac{\partial x^w}{\partial x_i} + x_i \frac{\partial f}{\partial x_i} x^w = w_i x^w + x_i \frac{\partial f}{\partial x_i} x^w. \tag{2}$$

There is a natural map

$$\phi : \mathcal{L}' \longrightarrow H_{\text{dR}}^{n+1}(\mathbb{P}^{n+1} \setminus X), \quad x^w \mapsto (-1)^{\gamma-1} (\gamma - 1)! \frac{x^w}{f^\gamma} \Omega^*.$$

Now we use the Poincaré residue map

$$\text{Res} : H_{\text{dR}}^{n+1}(\mathbb{P}^{n+1} \setminus X) \longrightarrow H_{\text{dR}}^n(X)$$

and observe that the composition $\rho := \text{Res} \cdot \phi$ factors through the quotient

$$W' = \mathcal{L}' / \mathcal{L}' \cap \sum_{i=0}^{n+1} D_i \mathcal{L}.$$

and we get an isomorphism $W' \cong H_{\text{dR}}^n(X)_0$ which is compatible with the pole order filtration on the source and the Hodge filtration on the target, cf. [8], cf. also [1],[15]. The image of $H^n(X, O_X)$ under the action of absolute Frobenius on $H_{\text{dR}}^n(X)$ is up to a multiplicative constant given by raising x^w/f^γ to the p -th power and then reducing the pole order by making use of the relations (2).

8. Fermat Varieties

We give examples of surfaces with a -number equal to 0, 1 or 2.

THEOREM 8.1. *Let p be a prime different from 5. Let X be the Fermat surface in characteristic p defined by $\sum_{i=0}^3 x_i^5 = 0$ in \mathbb{P}^3 . Then the a -number of X satisfies*

$$a(X) = \begin{cases} 0 & p \equiv 1 \pmod{5} \\ 1 & p \equiv 2, 3 \pmod{5} \\ 2 & p \equiv 4 \pmod{5}. \end{cases}$$

PROOF. We use the description of the (primitive) cohomology $H^2(X, O_X)$ with the Poincaré residue map

$$P : V \longrightarrow H_{\text{dR}}^2(X), \quad A \mapsto \text{Res}\left(\frac{A x_0 x_1 x_2 x_3}{f^3} \Omega^*\right)$$

with V the vector space of homogeneous polynomials of degree 11. A basis of the 4-dimensional space $H^2(X, O_X)$ is given by $\alpha_i = P((x_0x_1x_2x_3)^3/x_i)$ for $i = 0, \dots, 3$. Since Ω^* is a logarithmic form the image of α_0 under Frobenius is given by the Poincaré residue of

$$\frac{(x_0x_1x_2x_3)^{3p}(x_1x_2x_3)^p}{f^{3p}} \Omega^*$$

and using the relations $-5x_i^5x^w = w_ix^w$ for any monomial x^w obtained from (2) this is equivalent to an expression $\text{res}(g\Omega^*)$ with g given by

$$\begin{aligned} (x_0x_1x_2x_3)^3x_1x_2x_3/f^3 & p \equiv 1 \pmod{5} \\ (x_0x_1x_2x_3)(x_1x_2x_3)^2/f^2 & p \equiv 2 \pmod{5} \\ x_0^4(x_1x_2x_3)^2/f^2 & p \equiv 3 \pmod{5} \\ x_0^2x_1x_2x_3/f & p \equiv 4 \pmod{5} \end{aligned}$$

and similarly for the other α_i . This pole order implies that $a(X)$ is as indicated. □

THEOREM 8.2. *Let X be the Calabi-Yau Fermat variety in \mathbb{P}^r given by the equation $x_0^{r+1} + \dots + x_r^{r+1} = 0$ with $r + 1 \not\equiv 0 \pmod{p}$. Then the a -number of X is the natural number a with $0 \leq a \leq r - 1$ and $a \equiv p - 1 \pmod{r + 1}$.*

PROOF. Under the Poincaré residue map $(x_0 \dots x_r)^r \Omega^* / f^r$ maps to a generator of $H^{r-1}(X, O_X)$. The image under Frobenius is $(x_0 \dots x_r)^{pr} / f^{pr} \times \Omega^*$ and using again the relations (2) we can reduce the pole order pr modulo $r + 1$. But $pr \equiv -p \pmod{r + 1}$ and from this the result easily follows. □

There exist varieties of arbitrary dimension with maximal a -number namely abelian varieties which are products of supersingular elliptic curves. Besides these abelian varieties the following Calabi-Yau varieties give examples of such varieties.

COROLLARY 8.3. *Let X be the $(p - 1)$ -dimensional Fermat variety defined by*

$$x_0^{p+1} + x_1^{p+1} + \dots + x_p^{p+1} = 0$$

in projective space of dimension p . Then the a -number of the Calabi-Yau variety X is equal to $p - 1 = \dim(X)$.

Let X be the 7-dimensional cubic in \mathbb{P}^8 defined by $\sum_{i=0}^8 x_i^3 = 0$ in \mathbb{P}^8 . Then $H^7(X, O_X)$ and $H^6(X, \Omega_X^1)$ vanish, while $H^5(X, \Omega_X^2)$ is 1-dimensional. Let $a'(X)$ be the maximum j such that the Hodge step F^j contains G_2 (which can be seen as an image of $H^5(X, \Omega^2)$ under Frobenius divided by p^2 in $H_{\text{dR}}^2(X)$, cf. [7]).

THEOREM 8.4. *Let X be the 7-dimensional cubic in \mathbb{P}^8 defined by $\sum_{i=0}^8 x_i^3 = 0$. Then we have:*

$$a'(X) = \begin{cases} 2 & p \equiv 1 \pmod{3} \\ 5 & p \equiv 2 \pmod{3}. \end{cases}$$

PROOF. Using again the description of primitive cohomology with the Poincaré residue map we consider the form

$$\frac{(x_0 \dots x_8)^2}{f^6} \Omega^*$$

and then see that the image under Frobenius is equivalent to the Poincaré residue of $g\Omega^*$ with g given by $(x_0 \dots x_8)^2 / f^6$ if $p \equiv 1 \pmod{3}$ and $(x_0 \dots x_8) / f^3$ if $p \equiv 2 \pmod{3}$. □

9. Calabi-Yau Varieties

Let X be an n -dimensional Calabi-Yau variety. Such a variety has a height which we denote by $h = h(X)$. There are several possible definitions of this, e.g. with formal groups. Here we take as definition

$$h(X) - 1 := \max\{\dim H^{n-1}(X, B_i\Omega_X^1) : i \in \mathbb{Z}_{\geq 0}\}.$$

One can use the methods of [5] to see that this definition agrees with the definition using the formal group Φ^n associated to X . The natural inclusion $B_i\Omega_X^1 \subseteq \Omega_X^1$ induces a natural map $\beta_i : H^{n-1}(X, B_i\Omega_X^1) \rightarrow H^{n-1}(X, \Omega_X^1)$.

LEMMA 9.1. *If the natural map $\beta_i : H^{n-1}(X, B_i\Omega^1) \rightarrow H^{n-1}(X, \Omega^1)$ is not injective, then $\lim_{i \rightarrow \infty} \dim H^{n-1}(X, B_i\Omega_X^1) = \infty$.*

PROOF. The argument is similar to the cases of K3 surfaces, cf. [5]. We give the argument for $i = 1$. Suppose that β_1 has a non-trivial kernel represented by a cocycle $\{df_{\alpha_1, \dots, \alpha_n}\}$. Then we have a relation

$$df_{\alpha_1, \dots, \alpha_n} = \sum (-1)^j \omega_{\alpha_1, \dots, \hat{\alpha}_j, \dots, \alpha_n}.$$

Since the Cartier map $H^0(U, \Omega_{U, \text{closed}}^1) \rightarrow H^0(U, \Omega_U^1)$ is surjective on affine sets U we can find closed forms $\tilde{\omega}_{\alpha_1, \dots, \hat{\alpha}_j, \dots, \alpha_n}$ and regular functions $g_{\alpha_1, \dots, \alpha_n}$ on $\cap U_{\alpha_i}$ and obtain a new relation

$$f_{\alpha_1, \dots, \alpha_n}^{p-1} df_{\alpha_1, \dots, \alpha_n} + dg_{\alpha_1, \dots, \alpha_n} = \sum (-1)^j \tilde{\omega}_{\alpha_1, \dots, \hat{\alpha}_j, \dots, \alpha_n}.$$

One now checks directly that the cocycle at the left hand side represents an element of $H^{n-1}(X, B_2\Omega_X^1)$ which does not lie in the image of $H^{n-1}(X, B_1\Omega_X^1)$. One also checks that this element is non-zero. The argument for other i is similar. \square

PROPOSITION 9.2. *If $h \neq \infty$ then $1 \leq h \leq h^{1, n-1} + 1$.*

PROOF. If $h \neq \infty$ all the maps β_i are injective. \square

REMARK 9.3. We have $h^1(\Theta) = h^1(\Omega^{n-1})$, so we a priori could expect a stratification of the moduli with h steps; but the example of K3 surfaces shows that h cannot assume all values between 1 and $h^{1,1} + 1$. Therefore one should find bounds for h .

PROPOSITION 9.4. *Let X be a Calabi-Yau variety such that the Hodge-to-de Rham spectral sequence degenerates. Then we have either $h(X) = 1$ and $a(X) = 0$, or $2 \leq h(X) < \infty$ and $a(X) = 1$, or $h(X) = \infty$ and $a(X) \geq 1$.*

PROOF. We have $h(X) = 1$ if and only if the map $F_{\text{abs}} : H^n(X, O_X) \rightarrow F^0/F^1 = H^n(X, O_X)$ is non-zero and this is equivalent to $F_{\text{abs}}(H^n(X, O_X)) \not\subseteq F^1$. So we have $h(X) \geq 2$ if and only if the map F maps $H^n(X, O_X)$ to F^1 and then we have the projection $H^n(X, O_X) \rightarrow F^1 \rightarrow F^1/F^2 = H^{n-1}(X, \Omega_X^1)$. Now recall that we have an isomorphism $d : O_X/FO_X \cong B_1\Omega_X^1$ and from the exact sequence

$$0 \rightarrow O_X \xrightarrow{F} O_X \rightarrow O_X/FO_X \rightarrow 0$$

we thus get $H^{n-1}(X, B_1\Omega_X^1) \cong H^n(X, O_X)$. We claim that the image of $H^n(X, O_X)$ under F in F^1/F^2 is the image of $H^{n-1}(X, B_1\Omega_X^1)$ in $H^{n-1}(X, \Omega_X^1)$. This can be checked by a direct computation: if $\{f_I\}$ is a Čech n -cocycle representing a generator g of $H^n(X, O_X)$ then $F(g)$ is represented by $\{f_I^p\}$ and this is cohomologous to 0 in $H^n(X, O_X)$. So there exists an $(n-1)$ -cocycle h_J such that $\delta(h) = f^p$. The

image of g in F^1/F^2 is represented by dh_J . Now use the proof of the preceding Proposition. \square

COROLLARY 9.5. *Let X be the Fermat Calabi-Yau variety of degree $r+1$ in \mathbb{P}^r . Assume that the characteristic p does not divide $r+1$ and satisfies $p \not\equiv 2 \pmod{r+1}$. Then the height $h(X)$ is equal to 1 or ∞ . Moreover, $h(X) = 1$ if and only if $p \equiv 1 \pmod{r+1}$.*

REMARK 9.6. The corollary holds also without the condition that $p \not\equiv 2 \pmod{r+1}$, but for the proof we then need to use Jacobi sums.

10. Relation with a Result of Ogus

In [12] Ogus proved a result on the order of vanishing of the Hasse-invariant of a family of Calabi-Yau varieties that is closely related to the a -number introduced here. Let $f : X \rightarrow S$ be a family of Calabi-Yau varieties such that for each fibre X_s the Hodge-to-de Rham spectral sequence degenerates and such that the Kodaira-Spencer mapping $T_{S/k} \rightarrow R^1 f_*(T_{X/S})$ is surjective. Then Ogus proves that the Hasse-invariant vanishes to order i at s if $G_0(X_s) \subset F^i(X_s)$. Moreover, under an additional natural assumption on X/S the order of vanishing is exactly equal to the a -number of X_s .

As an example we consider the family of Calabi-Yau hypersurfaces in \mathbb{P}^4 given by

$$X_\alpha : \quad \sum x_i^5 - 5\alpha x_1 \cdots x_5 = 0.$$

We have $h^{1,1} = 1$, $h^{2,1} = 101$ and $b_3 = 204$. The Hasse-Witt invariant for $p \neq 5$ can be calculated (cf. [7], §2.3)

$$H(\alpha) = \sum_{m=0}^{\lfloor (p-1)/5 \rfloor} \frac{(5m!)}{(m!)^5} \alpha^{p-1-5m}.$$

This is a polynomial of degree $p-1$ in α . For $\alpha = 0$ we find by using Theorem 8.2 and the formula for $H(\alpha)$:

PROPOSITION 10.1. *Let p be a prime $\neq 5$. Then the a -number of X_0 is determined by*

$$a(X_0) = \text{ord}_0 H(\alpha) \equiv p-1 \pmod{5}.$$

We have an action of the symmetric group S_5 by permutation of the coordinates and of μ_5^3 given by generators

$$\begin{aligned} & \xrightarrow{g_1} (x_1, \zeta x_2, x_3, x_4, \zeta^4 x_5) \\ (x_1, x_2, x_3, x_4, x_5) & \xrightarrow{g_2} (x_1, x_2, \zeta x_3, x_4, \zeta^4 x_5) \\ & \xrightarrow{g_3} (x_1, x_2, x_3, \zeta x_4, \zeta^4 x_5) \end{aligned}$$

Then $Y_\alpha = X_\alpha/\mu_5^3$ is a mirror family ([11], [2]) with $h^{1,1} = 101$, $h^{2,1} = 1$ and $b_3 = 4$. We see a 4-step filtration

$$F^3 \subset F^2 \subset F^1 \subset H_{\text{dR}}^3.$$

It is not difficult to see, with the help of a non-trivial trace map $H^i(X_\alpha, \Omega_{X_\alpha}^j) \rightarrow H^i(Y_\alpha, \Omega_{Y_\alpha}^j)$, that the a -number of Y_α equals that of X_α .

11. Acknowledgements

This research was made possible by a JSPS-NWO grant. The second author would like to thank the University of Amsterdam and the first author would like to thank Prof. Ueno for inviting him to Kyoto, where this paper was finished. We also thank Ben Moonen for his comments.

References

- [1] J. Carlson, Ph. Griffiths: Infinitesimal variations of Hodge structure and the global Torelli problem. In: *Journées de Géométrie Algébrique d'Angers, 1979*, pp. 51–76, Sijthoff & Noordhoff, Alphen aan den Rijn—Germantown, Md., 1980.
- [2] P. Deligne: Local behavior of Hodge structures at infinity. *Mirror symmetry, II*, 683–699, AMS/IP Stud. Adv. Math., 1, Amer. Math. Soc., Providence, RI, 1997.
- [3] T. Ekedahl, G. van der Geer: In preparation.
- [4] G. van der Geer: Cycles on the moduli space of abelian varieties. In: *Moduli of curves and abelian varieties*, 65–89, Aspects Math., E33, Vieweg, Braunschweig, 1999.
- [5] G. van der Geer, T. Katsura: On a stratification of the moduli of $K3$ surfaces. *J. Eur. Math. Soc.* 2 (2000), 259–290.
- [6] L. Illusie, M. Raynaud: Les suites spectrales associées au complexe de de Rham-Witt. *Inst. Hautes Études Sci. Publ. Math.* 57 (1983), 73–212.
- [7] N. Katz: Algebraic solutions of differential equations (p -curvature and the Hodge filtration). *Invent. Math.* 18 (1972), 1–118.
- [8] N. Katz: On the intersection matrix of a hypersurface. *Ann. Sci. École Norm. Sup.* (4) (1969) 583–598.
- [9] N. Koblitz: p -adic variation of the zeta-function over families of varieties defined over finite fields. *Compositio Math.* 31 (1975), 119–218.
- [10] B. Moonen: Group schemes with additional structures and Weyl group cosets. In *Moduli of abelian varieties (Texel Island, 1999)*, 255–298, Progr. Math., 195, Birkhäuser, Basel, 2001.
- [11] D. Morrison: Mirror symmetry and rational curves on quintic threefolds: a guide for mathematicians. *J. Amer. Math. Soc.* 6 (1993), 223–247.
- [12] A. Ogus: On the Hasse locus of a Calabi-Yau family. *Math. Res. Lett.* 8 (2001), 35–41.
- [13] F. Oort: Commutative group schemes. *Lecture Notes in Mathematics*, 15 Springer-Verlag, Berlin-New York 1966.
- [14] F. Oort: A stratification of a moduli space of abelian varieties. In *Moduli of abelian varieties (Texel Island, 1999)*, 345–416, Progr. Math., 195, Birkhäuser, Basel, 2001.
- [15] J. Steenbrink: Intersection form for quasi-homogeneous singularities. *Compositio Math.* 34 (1977), no. 2, 211–223.

FACULTEIT WISKUNDE EN INFORMATICA, UNIVERSITY OF AMSTERDAM, PLANTAGE MUIDERGRACHT 24, 1018 TV AMSTERDAM, THE NETHERLANDS
E-mail address: `geer@science.uva.nl`

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, KOMABA, MEGURO-KU, TOKYO, 153-8914 JAPAN
E-mail address: `tkatsura@ms.u-tokyo.ac.jp`